



MAX-PLANCK-GESELLSCHAFT

# Search for supersymmetry in final states with jets and missing transverse momentum with the ATLAS detector

Michael Flowerdew, Max Planck  
Institute for physics, Munich  
on behalf of the ATLAS  
collaboration

Supersymmetry 2011, Fermilab

# Introduction

- Searches for new **strongly interacting** particles in final states with **jets and missing transverse momentum**
- $>1 \text{ fb}^{-1}$  analysed at  $\sqrt{s} = 7 \text{ TeV}$

## 2-4 jet inclusive search


$$\tilde{q}\tilde{q}, \tilde{q}\tilde{g}, \tilde{g}\tilde{g}$$

$$\tilde{q} \rightarrow q\tilde{\chi}_1^0$$

$$\tilde{g} \rightarrow q\bar{q}\tilde{\chi}_1^0$$

## 6-8 jet inclusive search

Many-body decays  
Non-leptonic  
cascade decays



# Object selection

\*For veto:  
Tighter selections  
used in leptonic  
Control Regions  
(eg  $p_T > 25$  GeV)

## Jets

Anit- $k_T$  algorithm with  $R=0.4$   
 $p_T > 20$  GeV,  $|\eta| < 2.8$   
MC-based calibration with pile-up and  
vertex corrections

## Missing transverse momentum (MET)

Jet-based MET + out-of-cluster  
contributions  
Corrections for loosely-selected  
electrons and muons ( $p_T > 10$  GeV)

## Trigger

>95% efficient in signal regions

## Electrons and muons\*

$p_T > 20$  GeV,  $|\eta_e| < 2.47$ ,  $|\eta_\mu| < 2.40$   
Muon isolation:  $\Sigma p_T (\Delta R < 0.2) < 1.8$  GeV

## Overlap removal

$\Delta R(e, \text{jet}) < 0.2 \Rightarrow$  remove jet  
 $\Delta R(e/\mu, \text{jet}) < 0.4 \Rightarrow$  remove lepton

## Event veto

Primary vertex has  $< 5$  tracks  
Jet quality + other event cleaning  
- Including LAr readout problems;  
veto region of size  $1.4 \times 0.2$  in  $\Delta\eta \times \Delta\phi$   
Reconstructed electron or muon  
- covered by independent analyses

# 2-4 JET ANALYSIS



# Signal Region selection

	Signal Region	$\geq 2$ jets	$\geq 3$ jets	$\geq 4$ jets	High mass
Trigger plateau	$E_T^{\text{miss}}$	$> 130$	$> 130$	$> 130$	$> 130$
	Leading jet $p_T$	$> 130$	$> 130$	$> 130$	$> 130$
Jet multiplicity	Second jet $p_T$	$> 40$	$> 40$	$> 40$	$> 80$
	Third jet $p_T$	–	$> 40$	$> 40$	$> 80$
	Fourth jet $p_T$	–	–	$> 40$	$> 80$
QCD rejection	$\Delta\phi(\text{jet}, E_T^{\text{miss}})_{\text{min}}$	$> 0.4$	$> 0.4$	$> 0.4$	$> 0.4$
Signal definition	$E_T^{\text{miss}}/m_{\text{eff}}$	$> 0.3$	$> 0.25$	$> 0.25$	$> 0.2$
	$m_{\text{eff}}$ [GeV]	$> 1000$	$> 1000$	$> 500/1000$	$> 1100$

$\tilde{q}\tilde{q}$        $\tilde{q}\tilde{g}$        $\tilde{g}\tilde{g}$

$\int L dt = 1.04 \text{ fb}^{-1} \pm 0.04 \text{ fb}^{-1}$   
 $m_{\text{eff}} = E_T^{\text{miss}} + \sum_{\text{SR jets}} p_T$   
 $m_{\text{eff}}^{\text{incl}} = E_T^{\text{miss}} + \sum_{p_T > 40 \text{ GeV}} p_T^{\text{jet}}$

- Jet + MET trigger
- 5 Signal Regions, targeting different topologies and mass ranges
- Optimised for discovery using a simplified SUSY model


# Analysis strategy

- Principal background components:
  - **W/Z+jets** ( $Z \rightarrow \nu\nu$  irreducible)
  - **Top**
  - QCD multi-jets
- Estimated using **Control Regions**, five for each Signal Region
  - Input to combined likelihood fit
- **Transfer Factors** relate CR measurement to SR background estimate
  - Ratio reduces some systematic uncertainties

$$N(\text{SR}, \text{est}, \text{proc}) = N(\text{CR}, \text{obs}, \text{proc}) * \left[ \frac{N(\text{SR}, \text{raw}, \text{proc})}{N(\text{CR}, \text{raw}, \text{proc})} \right]$$

- **Profile likelihood fit** accounts for correlated systematic uncertainties and CR contamination


# QCD background




- **Data-driven background estimation**
  - Control region defined by  $\Delta\phi(\text{jet}, \text{MET})_{\min} < 0.2$ 
    - Mis-measured jets
    - Heavy flavour
  - Transfer Factor computed by **smearing jets** in low-MET events
    - Smearing produces high MET events => count SR/CR ratio using  $\Delta\phi$
  - Special treatment of region with **LAr readout problems**
- Uncertainties from modelling of jet smearing

# Z+jets background

- Two Control Regions
  - $\gamma$ +jets, with photon treated as MET
  - $Z(\rightarrow ll)$ +jets, with Z treated as MET
- Transfer Factors taken from simulation
- Main uncertainties:
  - Theoretical extrapolations
  - Jet energy scale/resolution
  - Other detector systematics
  - MC statistics




# W and top background



- CR events selected with a **lepton + MET** selection
  - $30 < m_T < 100 \text{ GeV}^*$
  - **b-tagged jet**  $\Rightarrow$  top CR
  - Otherwise  $\Rightarrow$  W CR
  - Lepton treated as jet for other kinematic cuts
- Transfer Factor from simulation
- Main uncertainties:
  - Theoretical extrapolations
  - Jet energy scale/resolution
  - Other detector systematics
  - Pile-up
  - b-tagging uncertainties
  - MC statistics

$$^* m_T = \sqrt{2p_T^\ell E_T^{\text{miss}} [1 - \cos \Delta\phi(p_T^\ell, E_T^{\text{miss}})]}$$

# Signal Region observations



- Data and raw MC agree well
- Model limits obtained using  $\text{CL}_s$  prescription



# Results

value  $\pm$  stat  $\pm$  syst

Process	Signal Region				
	$\geq 2$ -jet	$\geq 3$ -jet	$\geq 4$ -jet, $m_{\text{eff}} > 500$ GeV	$\geq 4$ -jet, $m_{\text{eff}} > 1000$ GeV	High mass
$Z/\gamma$ +jets	$32.5 \pm 2.6 \pm 6.8$	$25.8 \pm 2.6 \pm 4.9$	$208 \pm 9 \pm 37$	$16.2 \pm 2.1 \pm 3.6$	$3.3 \pm 1.0 \pm 1.3$
$W$ +jets	$26.2 \pm 3.9 \pm 6.7$	$22.7 \pm 3.5 \pm 5.8$	$367 \pm 30 \pm 126$	$12.7 \pm 2.1 \pm 4.7$	$2.2 \pm 0.9 \pm 1.2$
$t\bar{t}$ + Single Top	$3.4 \pm 1.5 \pm 1.6$	$5.6 \pm 2.0 \pm 2.2$	$375 \pm 37 \pm 74$	$3.7 \pm 1.2 \pm 2.0$	$5.6 \pm 1.7 \pm 2.1$
QCD jets	$0.22 \pm 0.06 \pm 0.24$	$0.92 \pm 0.12 \pm 0.46$	$34 \pm 2 \pm 29$	$0.74 \pm 0.14 \pm 0.51$	$2.10 \pm 0.37 \pm 0.83$
Total	$62.3 \pm 4.3 \pm 9.2$	$55 \pm 3.8 \pm 7.3$	$984 \pm 39 \pm 145$	$33.4 \pm 2.9 \pm 6.3$	$13.2 \pm 1.9 \pm 2.6$
Data	58	59	1118	40	18


- Model-independent limit** on the *uncorrected* non-SM cross section within our observable signal regions

Process	Signal Region				
	$\geq 2$ -jet	$\geq 3$ -jet	$\geq 4$ -jet, $m_{\text{eff}} > 500$ GeV	$\geq 4$ -jet, $m_{\text{eff}} > 1000$ GeV	High mass
Excluded $\sigma \times \text{Acc} \times \epsilon$ (fb)	24	30	477	32	17




# Interpretation: Simple model

- **Three-particle model**
  - **Squarks** (two generations, with same mass)
  - **Gluinos**
  - **LSP** ( $m = 0$ )
  - $m = 5 \text{ TeV} \forall$  other sparticles
- „Perfect“ signature for this analysis
- Other  $m_{\text{LSP}}$  scenarios to be explored in upcoming paper
  - Low dependence up to  $\sim 200 \text{ GeV}$
- Channel with best expected limit chosen for each point
- **Exclusion extending up towards  $m \sim 1 \text{ TeV}$**




# Interpretation: CMSSM/ mSUGRA

- $A_0 = 0$
- $\mu > 0$
- $\tan \beta = 10$
- **Easy comparison to older results**
- **Extended reach in  $m_0$**  due to new signal regions
- $m_{1/2} > 450$  GeV at low  $m_0$
- Other model interpretations possible – follow up with information on HEPDATA



# 6-8 JET ANALYSIS

# Large jet multiplicity analysis



- Extension of 2-4 jet analysis
  - Increased sensitivity to **many-body** or **cascade decays**
  - Example: **high  $m_0$**  region in CMSSM/mSUGRA
- Signature: **>6 to >8 jets** + **MET** + lepton veto
  - QCD modelling is **the** issue
  - We cannot expect a good MC prediction => **Entirely data-driven approach**

$$\text{MET} / \sqrt{H_T}$$


(including fully hadronic tt)

- MET resolution dominated by **stochastic fluctuations**

$$\sigma^2(E_T^{\text{miss}}) \sim H_T = \sum p_T^{\text{jet}}$$

- Use  $\text{MET} / \sqrt{H_T}$  to **remove MC simulation dependence**
  - Nearly independent of jet multiplicity and pile-up
  - Cut on this, and **no other MET variables**

5 jet selection compared  
with 4 jet template




$$\text{MET}/\sqrt{H_T}$$

(including fully hadronic  $t\bar{t}$ )

- Control Region with  $\text{MET}/\sqrt{H_T} < 1.5 \sqrt{\text{GeV}}$
- Transfer Factor from events with exactly 5 or 6 jets
- Systematic cross-checks:**
  - $\text{MET}/\sqrt{H_T}$  invariance tested using lower-multiplicity selections
  - Jet smearing, as in 2-4 jet analysis
  - Jet flavour separation
  - Special study of LAr readout-problem region

5 jet selection compared with 4 jet template





# Signal Region selection

- **Multi-jet triggers**, requiring 4 or 5 jets
  - => Lower leading jet threshold
- **MET/ $\sqrt{H_T}$**  replaces MET and MET/ $m_{\text{eff}}$ 
  - $\Delta\phi(\text{jet}, \text{MET})_{\text{min}}$  also removed
- **LAr readout problem** => Jet energy correction in affected region, event vetoed if MET correction is large (>10 GeV and >10% of MET)

	Signal region	7j55	8j55	6j80	7j80
Trigger plateau	Jet $p_T$	> 55 GeV		> 80 GeV	
	Jet $ \eta $	< 2.8			
	$\Delta R_{jj}$	> 0.6 for any pair of jets			
SR definition	Number of jets	$\geq 7$	$\geq 8$	$\geq 6$	$\geq 7$
BG reduction and control	$E_T^{\text{miss}}/\sqrt{H_T}$	> 3.5 GeV <sup>1/2</sup>			

Trigger plateau

$$\int L dt = 1.34 \text{ fb}^{-1}$$

$$H_T = \sum_{\text{Jets}} p_T$$

( $p_T > 40 \text{ GeV}, |\eta| < 2.8$ )





# Top and other backgrounds

(Semi- and fully leptonic  $t\bar{t}$ , W/Z + jets)

- **Top:** (Second largest background)
  - **Control Region:** 1 muon,  $40 < m_T < 100$  GeV, b-tagged jet
    - SR jet cuts applied, treating muon as a jet
  - **Transfer Factors** from simulation (ALPGEN)
    - Systematic uncertainties estimated as in 2-4 jet analysis
  - **Validation Regions (VRs)** vary jet  $p_T$  and MET/ $\sqrt{H_T}$  cuts
- **W/Z+jets** are small:
  - Estimated from **simulation**, due to low CR statistics
  - Validation Regions show no problems
    - W: Like top, but b-jet veto
    - Z: Two muon VR selection

# Validation Regions

- Use 4 & 5 jet Validation Regions, extending out in  $N_{\text{jet}}$  as far as statistics allow



- Jet multiplicity tested in events with and without leptons
- Scaling of the number of jets is understood
- Also agreement with other theoretical cross-checks

# Results

Signal region	7j55	8j55	6j80	7j80
Multi-jets	$26 \pm 5.2$	$2.3 \pm 0.7$	$19 \pm 4$	$1.3 \pm 0.4$
$t\bar{t} \rightarrow \ell(\ell) X$	$10.8 \pm 6.7$	$0^{+4.3}$	$6.0 \pm 4.6$	$0^{+0.13}$
$W + \text{jets}$	$0.95 \pm 0.80$	$0^{+0.13}$	$0.34 \pm 0.34$	$0^{+0.13}$
$Z + \text{jets}$	$1.5^{+1.8}_{-1.5}$	$0^{+0.75}$	$0^{+0.75}$	$0^{+0.75}$
<b>Total SM</b>	$39.3^{+8.7}_{-8.5}$	$2.3^{+4.4}_{-0.7}$	$25.8 \pm 6.1$	$1.3^{+0.9}_{-0.4}$
<b>Data</b>	45	4	26	3
$N_{\text{BSM,max}}^{95\%}$	26.0	11.2	16.3	6.0
$(\sigma_{\text{BSM,max}}^{95\%} \times \epsilon)/\text{fb}$	19.4	8.4	12.2	4.5
$p_{\text{SM}}$	0.30	0.36	0.49	0.16


- No excess seen => set limits
- $N$  and  $\sigma$  are **model-independent upper limits** on non-SM processes after all selection
  - Still using  $\text{CL}_s$  prescription

# Example interpretation: CMSSM

Exclusion from  
2-4 jet analysis

- Reminder:  $A_0 = 0$ ,  $\mu > 0$ ,  $\tan \beta = 10$
- Channel with best expected limit chosen for each point
- Upward fluctuations in two key channels => exclusion less than predicted

$$m_{\tilde{g}} > 520 \text{ GeV}$$



# Conclusion

- The search reach in jets+MET final states has been **dramatically extended**
  - 2-4 jet and 6-8 jet inclusive multiplicities
  - $\sim 1 \text{ fb}^{-1}$  of data & analysis improvements
  - Negative results interpreted using simplified *R*-parity conserving SUSY model and CMSSM
  - Mass reach at or approaching 1 TeV
- Papers are in preparation for these analyses

If you want more...

**BACKUP**



# Statistical methods


- Simultaneous likelihood fit to Signal Region + 5 Control Regions in each channel
  - Six Poisson-distributed variables and PDF to constrain systematic uncertainties






$$L(n | \mu, b, \theta) = P_{\text{SR}} \times P_{\text{WR}} \times P_{\text{TR}} \times P_{\text{ZR}_a} \times P_{\text{ZR}_b} \times P_{\text{QR}} \times C_{\text{syst}}$$

- Correlations between Control Regions taken into account
  - eg jet energy scale and b-tagging efficiency








# Signal Region observations















# Dijet channel Control Regions



# Three jet channel Control Regions




# Four jet channel Control Regions




Four jet  
high mass  
channel  
Control  
Regions

# Dijet channel exclusion






# Three jet channel exclusion







# Four jet channel exclusion ( $m_{\text{eff}} > 500 \text{ GeV}$ )





# Four jet channel exclusion ( $m_{\text{eff}} > 1000 \text{ GeV}$ )





# Four jet high mass channel exclusion





# MET/ $\sqrt{H_T}$ validation



# Jet multiplicity:



$$1.5 < \text{MET} / \sqrt{H_T} < 2.0 \sqrt{\text{GeV}}$$


# Jet multiplicity:



$$2.0 < \text{MET} / \sqrt{H_T} < 3.0 \sqrt{\text{GeV}}$$






# Multijets: Top Validation Region





# Multijets: W Validation Region



# Multijets: Z Validation Region



# MET/ $\sqrt{H_T}$ in the Signal Regions



# 6-8 jet study exclusion vs 2010 2-4 jets results

